



Study of the effect of neutrino oscillation on the supernova neutrino signal with the LVD detector

M. Aglietta^a, P. Antonioli^b, G. Bari^b, C. Castagnoli^a, W. Fulgione^a, P. Galeotti^a, M. Garbini^b, P.L. Ghia^a, P. Giusti^b, E. Kemp^c, A.S. Malguin^d, H. Menghetti^b, A. Pesci^b, I.A. Pless^e, A. Porta^a, V.G. Rzasny^d, O.G. Ryazhskaya^d, O. Saavedra^a, G. Sartorelli^b, M. Selvi^b, C. Vigorito^a, F. Vissani^f, L. Votano^g, V.F. Yakushev^d, G.T. Zatsepin^d, A. Zichichi^b

^aInstitute of Physics of Interplanetary Space, IFSI, CNR, Torino, University of Torino and INFN-Torino, Italy,

^bUniversity of Bologna and INFN-Bologna, Italy,

^cUniversity of Campinas, Campinas, Brazil,

^dInstitute for Nuclear Research, Russian Academy of Sciences, Moscow, Russia,

^eMassachusetts Institute of Technology, Cambridge, USA,

^fINFN-LNGS, Assergi, Italy

^gINFN-LNF, Frascati, Italy

We present an update of our previous study [1] on how ν oscillations affect the signal from a supernova core collapse observed in the LVD detector at LNGS. In this paper we use a recent, more precise determination of the cross section [2] to calculate the expected number of inverse beta decay events; we introduce in the simulation also the ν -Fe interactions; we include the Earth matter effects; and finally we study also the inverted mass hierarchy case.

1. SUPERNOVA NEUTRINO SIGNAL

At the end of its burning phase, a massive star ($M \geq 8M_{\odot}$) explodes into a supernova (SN), originating a neutron star which cools, emitting about 99 % of the liberated gravitational binding energy in neutrinos. The time-integrated spectra can be well approximated by the pinched Fermi–Dirac distribution, with an effective degeneracy parameter η , which is assumed, for simplicity, to be null. Although the hierarchy $\langle E_{\nu_e} \rangle < \langle E_{\bar{\nu}_e} \rangle < \langle E_{\nu_x} \rangle$ remains valid (here ν_x refers to both ν_{μ} and ν_{τ}), recent studies, with an improved treatment of ν transport, microphysics, the inclusion of the nucleon bremsstrahlung, and the energy transfer by recoils, find somewhat smaller differences between the $\bar{\nu}_e$ and ν_x spectra [3].

In the following, we assume a future galactic

SN explosion at a typical distance of $D = 10$ kpc, with a binding energy of $E_b = 3 \cdot 10^{53}$ erg and a total energy of $E_{\nu_e}^{\text{tot}} = E_{\bar{\nu}_e}^{\text{tot}} = E_{\nu_x}^{\text{tot}}$. We also assume that the fluxes of ν_{μ} , ν_{τ} , $\bar{\nu}_{\mu}$, and $\bar{\nu}_{\tau}$ are identical; we fix $T_{\nu_x}/T_{\bar{\nu}_e} = 1.5$, $T_{\nu_e}/T_{\bar{\nu}_e} = 0.8$ and $T_{\bar{\nu}_e} = 5$ MeV [5].

2. LVD DETECTOR AND NEUTRINO REACTIONS

The Large Volume Detector (LVD) in the INFN Gran Sasso National Laboratory, Italy, consists of an array of 840 liquid scintillator (LS) counters, 1.5 m³ each. These are interleaved by streamer tubes, and arranged in a modular geometry; a detailed description is in [4]. The active scintillator mass is $M = 1000$ t. There are two subsets of counters: the external ones (43%), oper-




Figure 1. Cross sections of the main neutrino interactions in LVD.

ated at energy threshold $\mathcal{E}_h \simeq 7$ MeV; and inner ones (57%), better shielded from rock radioactivity and operated at $\mathcal{E}_h \simeq 4$ MeV.

In the following, we will focus on ν reactions with free protons and iron nuclei: (1) $\bar{\nu}_e p, e^+ n$, observed through a prompt signal from e^+ above threshold \mathcal{E}_h (detectable energy $E_d \simeq E_{\bar{\nu}_e} - 1.8$ MeV $+ 2m_e c^2$), followed by the signal from the $np, d\gamma$ capture ($E_\gamma = 2.2$ MeV), above \mathcal{E}_l and with a mean delay $\Delta t \simeq 180 \mu\text{s}$. The cross section for this reaction has been recently recalculated [2] with a better treatment of the 10 – 100 MeV region, i.e. the SN neutrino energy. The cross section behaviour with energy is shown in figure 1. The efficiency for the prompt signal is $\epsilon_{\bar{\nu}_e p, e^+ n} = 95\%$. The total number of free protons in the LS is $9.34 \cdot 10^{31}$.

The LVD detector presents an iron support structure made basically by two components: the tank (mean thickness: 0.4 cm) which contains the LS and the portatank (mean thickness: 1.5 cm) which hosts a cluster of 8 tanks. Indeed, the higher energy part of the ν flux could be detected also with the $\nu(\bar{\nu})\text{Fe}$ interaction, which results in an electron (positron) that could exit iron and release energy in the LS. The considered reactions are: (2) $\nu_e {}^{56}\text{Fe}, {}^{56}\text{Co} e^-$, the bind-




Figure 2. Effect of neutrino oscillations in the signal detected in LVD.

ing energy difference between the ground levels is $E_b^{\text{Co}} - E_b^{\text{Fe}} = 4.566$ MeV; moreover the first Co allowed state is at 3.589 MeV. Indeed, in this work, we considered $E_{e^-} = E_{\nu_e} - 8.15$ MeV. A full simulation of the LVD support structure and LS geometry has been developed in order to get the efficiency for an electron, generated randomly in the iron structure, to reach the LS with energy higher than \mathcal{E}_h . It is greater than 20% for $E_\nu > 30$ MeV and grows up to 70% for $E_\nu > 100$ MeV. On average, the electron energy detectable in LS is $E_d \simeq 0.45 \times E_\nu$. The total number of iron nuclei is $7.63 \cdot 10^{30}$.

(3) $\bar{\nu}_e {}^{56}\text{Fe}, {}^{56}\text{Mn} e^+$, the energy threshold is very similar to reaction (2) and the same considerations could be done. The cross section for reactions (2), (3) are taken respectively from [5,6] and plotted in figure 1.

3. NEUTRINO OSCILLATION AND MSW EFFECT IN THE SN AND IN THE EARTH

In the study of SN neutrinos, ν_μ and ν_τ are indistinguishable because of the corresponding charged lepton production threshold; consequently, the relevant parameters are just

$(\Delta m_{\text{sol}}^2, U_{e2}^2)$ and $(\Delta m_{\text{atm}}^2, U_{e3}^2)$. We will adopt the following numerical values: $\Delta m_{\text{sol}}^2 = 7 \cdot 10^{-5} \text{eV}^2$, $\Delta m_{\text{atm}}^2 = 2.5 \cdot 10^{-3} \text{eV}^2$, $U_{e2}^2 = 0.33$; the selected solar parameters $(\Delta m_{\text{sol}}^2, U_{e2}^2)$ describe the LMA-I solution, as it results from a global analysis including solar, CHOOZ and KamLAND ν data [7]. For a normal mass hierarchy (NH) scheme, ν (not $\bar{\nu}$) cross two resonance layers: one at higher density (H), which corresponds to $\Delta m_{\text{atm}}^2, U_{e3}^2$, and the other at lower density (L), corresponding to $\Delta m_{\text{sol}}^2, U_{e2}^2$. For inverted mass hierarchy (IH), transitions at the higher density layer occur in the $\bar{\nu}$ sector.

Given the energy range of SN ν (up to ~ 100 MeV) and considering a star density profile $\rho \propto 1/r^3$, the adiabaticity condition is always satisfied at the L resonance for any LMA solution, while at the H resonance, this depends on the value of U_{e3}^2 . When $U_{e3}^2 \geq 5 \cdot 10^{-4}$, the conversion is completely adiabatic, meaning that the flip probability between two adjacent mass eigenstates is null ($P_h = 0$). In the adiabatic case and NH, the $\bar{\nu}_e$ produced in the SN core arrive at Earth as ν_1 , and they have a high ($U_{e1}^2 \simeq \cos^2 \theta_{12} \simeq 0.7$) probability to be detected as $\bar{\nu}_e$. On the other hand, the original $\bar{\nu}_x$ arrive at Earth as ν_2 and ν_3 and are detected as $\bar{\nu}_e$ with probability $U_{e2}^2 \simeq \sin^2 \theta_{12}$. Given the higher energy spectrum of $\bar{\nu}_x$ this configuration results in a larger number of interactions, with respect to the no-oscillation case, due to the increasing cross sections with energy. In the adiabatic-IH case the detected $\bar{\nu}_e$ completely come from the original $\bar{\nu}_x$ flux in the star, and the number of interaction is still greater, as shown in figure 2. The oscillations scheme can be summarized as: $F_e = P_h U_{e2}^2 F_e^0 + (1 - P_h U_{e2}^2) F_x^0$ and $F_{\bar{e}} = U_{e1}^2 F_{\bar{e}}^0 + U_{e2}^2 F_{\bar{x}}^0$ for normal hierarchy; $F_e = U_{e2}^2 F_e^0 + U_{e1}^2 F_x^0$ and $F_{\bar{e}} = P_h U_{e1}^2 F_{\bar{e}}^0 + (1 - P_h U_{e1}^2) F_{\bar{x}}^0$ for inverted hierarchy, where F_{any}^0 are the original neutrino fluxes in the star and F_{any} are the observed ν fluxes. One can notice that, in the antineutrino channel, the non adiabatic ($P_h = 1$), IH case, is equivalent to the NH case (which does not depend on adiabaticity). In figure 3 is shown the contribution of $(\nu_e + \bar{\nu}_e)$ Fe interactions in the total number of events. For the chosen SN and oscillation parameters, they are about 18% of the signal.




Figure 3. Impact of iron interactions in the global neutrino signal in LVD.

If we consider the effect of Earth in the neutrino path to the detector, we must replace, in the detected flux estimation, U_{ei}^2 with P_{ei} ($i = 1, 2$), the probability for the mass eigenstate ν_i to be detected as ν_e after path in the Earth [8], which depends on the solar oscillation parameters and on the travelled density profile through the Earth. Earth matter effect are more relevant in the ν than in the $\bar{\nu}$ channel, so the effect in reaction (1) is quite weak (it also depends on the rather high Δm_{sol}^2), but it could be detected if compared with a high statistic sample (i.e. SK) or if a larger number of events is available, i.e. a closer SN.

REFERENCES

1. M. Aglietta *et al.*, Nucl. Phys. B Proc. Suppl. 110 (2002) 410-413.
2. A. Strumia and F. Vissani, astro-ph/0302055.
3. M.T. Keil, G.G. Raffelt, and H.-T. Janka, astro-ph/0208035.
4. M. Aglietta *et al.*, Il Nuovo Cimento A 105 (1992) 1793.
5. E. Kolbe and K. Langanke, nucl-th/0003060.
6. J. Toivanen *et al.*, Nucl. Phys. A 694 (2001)

395-408.

7. G.L. Fogli *et al.*, [hep-ph/0212127](#).
8. C. Lunardini and A. Smirnov,
[hep-ph/0106149](#).