

# Prospects of the Higgs Sector at HL-LHC



Paolo Giacomelli (INFN Bologna)  
*What Next ?*, LTS1 workshop, La Biodola  
Thursday, May 22th, 2014



# LHC Physics program priorities

The discovery of a SM-like scalar boson at  $m_H \sim 125$  GeV defines the physics priorities

- With LHC 13/14 TeV data until  $\sim 2022$  ( $\sim 300 \text{ fb}^{-1}$ )
  - Measure SM-like scalar boson properties
    - mass,  $J^{PC}$
    - individual couplings with 5-15% precision
  - Search for new physics at a higher mass scale (new energy region)
    - SUSY
    - Exotics



# Higgs Physics at HL-LHC

- What can we do at **HL-LHC** in the Higgs sector?  
until  $\sim 2032$  ( $\sim 3000 \text{ fb}^{-1}$ )
  - Measure existing decay channels with the highest precision
  - Observe rare Higgs decays
    - $H \rightarrow \mu\mu$
    - $H \rightarrow Z\gamma$
    - $H \rightarrow cc$  (?)
  - Double Higgs production (Higgs self-coupling)
  - Vector boson scattering
  - Look for small deviations from SM predictions



# From 2012 to HL-LHC

- From 30 to 3000  $\text{fb}^{-1}$ : two orders of magnitude extrapolation in luminosity

To calculate physics projections at HL-LHC



**Similar trigger and reconstruction performances as in 2012**

**Need upgraded detectors to offset the much harsher LHC conditions and radiation damage**

**ATLAS and CMS have launched a comprehensive upgrade program**



# Higgs boson projections after LS1


## Approaches adopted for physics projections

- **ATLAS:** perform physics studies using fast simulation to mimic the beam effects on momentum and energy resolution, acceptance, identification and reconstruction efficiencies, fake rates, etc.
- **CMS:** assume that an upgraded detector will compensate the effects of the higher pile-up, using three different scenarios:
  - Scenario 1: all systematic uncertainties are kept unchanged with respect to those in current data analyses
  - Scenario 2: the theoretical uncertainties are scaled by a factor of  $1/2$ , while other systematical uncertainties are scaled by  $1/\sqrt{L}$
  - Scenario 3: set theoretical uncertainties to zero, leave other syst. uncertainties the same as in 2012

# Higgs signal strength with 300 fb<sup>-1</sup>

- Extrapolation by two orders of magnitude to higher luminosity
  - is subject to large uncertainties
  - scenarios 1 and 2 provide likely upper and lower bounds
- Experience at LEP and Tevatron indicates that scaling with  $1/\sqrt{L}$  is not unrealistic

CMS Projection




300 fb<sup>-1</sup>, 14TeV (Scenario 1)  
 300 fb<sup>-1</sup>, 14TeV (Scenario 1)

$$\mu = \sigma/\sigma_{SM}$$

With 300 fb<sup>-1</sup> the precision on the signal strength,  $\mu = \sigma/\sigma_{SM}$ , is expected to be 10-15% per channel

# Higgs signal strength with 3000 fb<sup>-1</sup>

CMS Projection



$$\mu = \sigma/\sigma_{\text{SM}}$$


L (fb)	H→γγ	H→WW	H→ZZ	H→bb	H→ττ	H→Zγ	H→μμ	H→inv.
300	[6,12]	[6,11]	[7,11]	[11,14]	[8,14]	[62,62]	[40,42]	[17,28]
3000	[4,8]	[4,7]	[4,7]	[5,7]	[5,8]	[20,24]	[20,24]	[6,17]

With 3000 fb<sup>-1</sup> the precision on  $\mu$  is expected to be 4-8% per channel

# Higgs boson couplings @300 fb<sup>-1</sup>

- Two scenarios:
  - Scenario 1:** same systematics as in 2012
  - Scenario 2:** theory systematics scaled by a factor 1/2, other systematics scaled by 1/√L

CMS Projection



300 fb<sup>-1</sup>, 14 TeV (Scenario 1)

300 fb<sup>-1</sup>, 14 TeV (Scenario 1)

With 300 fb<sup>-1</sup> the uncertainties on the Higgs couplings are expected in the range  $\sigma(\kappa_V) \sim 4-7\%$

$\sigma(\kappa_f) \sim 6-15\%$


300 fb<sup>-1</sup> 14 TeV, Scenario 1

300 fb<sup>-1</sup> 14 TeV, Scenario 2



# Higgs boson couplings @3000 fb<sup>-1</sup>


CMS Projection




L (fb)	K	K	K	K	K	K	K	K	K	BR
300	[5,7]	[4,6]	[4,6]	[6,8]	[10,13]	[14,15]	[6,8]	[41,41]	[23,23]	[14,18]
3000	[2,5]	[2,5]	[2,4]	[3,5]	[4,7]	[7,10]	[2,5]	[10,12]	[8,8]	[7,11]

- With 3000 fb<sup>-1</sup> the Higgs couplings can be determined with high precision (2-7%)

CMS Projection



CMS Projection




- Extrapolation by two orders of magnitude to higher luminosity
  - is subject to large uncertainties
- Results will become syst. limited due to theory uncertainties. We must encourage our theoretical friends to improve their calculations!

# Higgs couplings @3000 fb<sup>-1</sup>

**ATLAS** Simulation Preliminary


$\sqrt{s} = 14$  TeV:  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$



- With 3000 fb<sup>-1</sup> the couplings can be determined with high precision (up to **a few %**)

# Higgs coupling ratios vs. mass

## Mass-scaled coupling ratios vs. particle mass










- By LHC14@300, we'll have probed all 3rd generation fermion couplings to  $O(10-20\%)$
- $H \rightarrow \mu^+\mu^-$  gives us access to 2nd lepton generation, i.e. is the mass-generation mechanism same for all generations, for quarks and leptons?

G. Salam, A. Weiler


mass  $\propto$  coupling to Higgs ?

$$Br(H \rightarrow \mu^+\mu^-)_{SM} = 2.2 \cdot 10^{-4}$$







- The decay  $H \rightarrow \mu\mu$  can be observed with a significance of 5 sigma
  - measurement of the  $H\mu\mu$  coupling with a precision of  $\sim 10\%$







# $H \rightarrow Z\gamma$

- $\gamma Z$  like  $\gamma\gamma$  and  $gg$  loop induced, but sensitive to effects invisible in  $\gamma\gamma$  and  $gg$  (because of chiral couplings)
- In composite Higgs: Not protected by Goldstone symmetry, large  $\gamma Z$  while  $\gamma\gamma$  and  $gg$  small

G. Salam, A. Weiler







- Hcc coupling can still be 4-8 x SM

$$\mathcal{L} = c_c h \frac{m_c}{v} \bar{c}c + \dots$$

- In composite Higgs


$$c_c \simeq 1 + \mathcal{O}\left(\frac{v^2}{f^2}\right) + \mathcal{O}\left(\epsilon_c^2 \frac{g_\psi^2 v^2}{m_\psi^2}\right)$$

large for composite  
charm and light charm  
partners

## Measuring it?

Like H → bb, but with  
charm tagging?


Or via H → J/ψ γ ? 1306.5770



G. Salam, A. Weiler

# Higgs boson pair-production

Destructive interference between the two diagrams



Many channels to investigate  
Most promising ones:

$b\bar{b}W^+W^-$  (large BR but large bkg.)

$b\bar{b}\gamma\gamma$  (clean but small BR)


$b\bar{b}\tau^+\tau^-$

$b\bar{b}\mu^+\mu^-$  also being considered

$b\bar{b}b\bar{b}$

$b\bar{b}ZZ \rightarrow b\bar{b}2l2\nu$

Taken from “Higgs self-coupling measurements at the LHC” by M. J.



NNLO cross-section at  $m_H=125$  GeV:

$$\sigma = 40 \pm 3 \text{ fb}$$

G. de Florian, J. Mazzitelli, [1309.6594](#)

# di-Higgs production

At HL-LHC with  $L=3000 \text{ fb}^{-1}$  we will produce  **$\sim 120000$**  HH events


However we pay a big price  
in BR's ...

$b\bar{b}W^+W^-$   $\sim 14000$  events

$b\bar{b}\gamma\gamma$   $\sim 150$  events



$b\bar{b}\tau^+\tau^-$   $\sim 4300$  events

$b\bar{b}2l2\nu$   $\sim 730$  events



# di-Higgs production

$$HH \rightarrow b\bar{b}\gamma\gamma$$



Preliminary results with  $L=3000 \text{ fb}^{-1}$



# di-Higgs production

$b\bar{b}W^+W^-$  is being studied. Looks very difficult

$b\bar{b}\tau^+\tau^-$  seems more promising, studies just began

$b\bar{b}2l2\nu$  could be an interesting possibility, studies not yet started

Higgs boson pair-production is a flagship channel of HL-LHC.

There is ongoing work in both experiments in order to be able to assess the full potential at HL-LHC.

## Personal opinion

There is good hope to reach a sensitivity of  $\sim 3\sigma$  per experiment with  $L=3000 \text{ fb}^{-1}$





# Conclusions



- ATLAS and CMS have exceeded their design performances during the first LHC run, showing that precision physics can be made under these conditions.
- The experience gained and a sound program of upgrades gives us confidence that the experiments will meet the physics expected at HL-LHC with  $3000 \text{ fb}^{-1}$ , collected at  $\sqrt{s}=14 \text{ TeV}$  and instantaneous luminosities of  $5 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ .
- Precision Higgs boson physics at HL-LHC is an attractive future scenario deserving substantial studies and R&D
  - it is a challenging project involving major upgrades of full detectors
  - Higgs boson couplings can be measured with few percent precision
  - rare Higgs boson decays can be probed
  - Higgs self-coupling studies possible
  - VV scattering will be probed
- LHC has an exciting physics program for the next twenty years!

# Backup





# Integrated luminosity in 2012

Integrated luminosity recorded in 2012:  $\sim 22 \text{ fb}^{-1}$

2011:  $L \sim 6 \text{ fb}^{-1}$

CMS Integrated Luminosity, pp, 2012,  $\sqrt{s} = 8 \text{ TeV}$

Data included from 2012-04-04 22:37 to 2012-12-16 20:49 UTC





Total delivered luminosity:  $\sim 30 \text{ fb}^{-1}$

Total recorded luminosity:  $\sim 27 \text{ fb}^{-1}$

Excellent LHC performance and very high data-taking efficiency of the two detectors


# New boson with a mass of $\sim 125$ GeV

- We have discovered a SM-like scalar boson with a mass of  $\sim 125$  GeV.
- $J^{PC}$ , consistent with SM scalar boson, couplings will need more data.



The new boson is consistent with being the SM Higgs boson

$$\mu = \sigma/\sigma_{SM} = 0.80 \pm 0.14$$





# LHC and HL-LHC




## LHC

Energy increase  
8 TeV to 13/14 TeV

Injection  
upgrade

## HL-LHC



Interaction  
region  
upgrade

$L_{instantaneous}$   
 $L_{integrated}$   
Pile Up

$8 \times 10^{33} Hz/cm^2$   
 $30 fb^{-1}$   
 $PU \sim 40$

$2 \times 10^{34} Hz/cm^2$   
 $300 fb^{-1}$   
 $PU \sim 50$

$5 \times 10^{34} Hz/cm^2$   
 $3000 fb^{-1}$   
 $PU \sim 140$

LS1

LS3

Phase 1 Upgrade

Phase 2 Upgrade

ATLAS, CMS  
Upgrade plan






# Detector and trigger challenges

- Need detectors and trigger with high performances from low to high energy scales
  - 125 GeV SM-like boson measurements
  - Multi-TeV new physics searches
- **Phase 1 Upgrade:** twice LHC design luminosity
  - Event pileup reaches ~50 collisions per beam crossing (@ 25 ns)
  - Factor 5 increase in trigger rates relative to 2012 run
- **Phase 2 Upgrade:** 5x LHC design luminosity
  - Event pileup reaches ~140 collisions per beam crossing (@ 25 ns)
  - Need solutions to cope with very high rates (10-15 x 2012), radiation and pileup


ATLAS and CMS were designed to cope with  $L = 1\text{-}2 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$

# Pileup in 2012



**Peak: 37 pileup events**

Design value  
**25 pileup events**  
( $L=10^{34}$ , BX=25 ns)





Basically, life will not be easy...

*Pileup at 25 ns and  $L = 2 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$*





# Upgrade challenges and recipe

Maintain low trigger thresholds, efficient particle and physics object reconstruction at high rate and pile-up


Need new technology R&Ds to:

- Increase granularity
- Increase data bandwidth
- Increase processing power
- Improve radiation hardness
- Minimize material in tracking devices



## LS1 Projects

- Complete Muon coverage (ME,RE4)
- Improve muon operation, DT electronics
- Replace HCAL photo-detectors in Forward (new PMTs) and Outer (HPD→SiPMs)
- DAQ1→DAQ2



LS1 (now)

LS2 (2018)

LS3 (~2023)

## Phase 1 Upgrades



- New Pixel detector, HCAL electronics and L1-Trigger upgrade
- GEMs for forward muon det. under review
- Preparatory work during LS1
  - New beam pipe for pixel upgrade
  - Install test slices of pixel, HCAL, L1-trigger
  - Install ECAL optical splitters for L1-trigger

## Phase 2: being defined now

- Tracker replacement, L1 Track-Trigger
- Forward: calorimetry, muons and tracking
- High precision timing for PU mitigation
- Further Trigger upgrade
- Further DAQ upgrade




# Expected Phase 1 improvements



Significant gain in signal reconstruction efficiency:

$H \rightarrow 4\mu$	+41%
$H \rightarrow 2\mu 2e$	+48%
$H \rightarrow 4e$	+51%




Total efficiency improvement:  
factor of 2.5 (4.5%  $\rightarrow$  11%)


Improved jet and MET  $\rightarrow$   
25% improvement in  $m_{\tau\tau}$   
resolution

# CMS Phase II Muon detector

Increase det. acceptance up to  $|\eta|=4.0$












# ATLAS upgrade program




## ATLAS has devised a 3 stage upgrade program

- |   |  |  |
|---|--|--|
| <ul style="list-style-type: none"> <li>• New insertable pixel b-layer (IBL)</li> <li>• New Al beam pipe</li> <li>• New pixel services</li> <li>• Complete installation of EE muon chambers</li> <li>• New evaporative cooling plant</li> <li>• Consolidation of detector services</li> <li>• Specific neutron shielding</li> <li>• Upgrade magnet cryogenics</li> </ul> | <ul style="list-style-type: none"> <li>• New Small Wheel (nSW) for the forward muon Spectrometer</li> <li>• High Precision Calorimeter L1-Trigger</li> <li>• Fast TrackIng (FTK) for L2-trigger</li> <li>• Topological L1-trigger processors</li> <li>• New forward diffractive physics detectors (AFP)</li> </ul> | <ul style="list-style-type: none"> <li>• Completely new tracking detector</li> <li>• Calorimeter electronics upgrades</li> <li>• Upgrade part of the muon system</li> <li>• Possible L1-trigger track trigger</li> <li>• Possible changes to the forward calorimeters</li> </ul> |
|---|--|--|





**di-photon mass  
resolution  
in  $t\bar{t}H$  channel**





# New LHC schedule

## LHC schedule beyond LS1

CMS Pixel installation

CMS target for LS3

Only EYETS (19 weeks) (no Linac4 connection during Run2)

LS2 starting in 2018 (July) 18 months + 3 months BC (Beam Commissioning)


LS3 LHC: starting in 2023 => 30 months + 3 BC

injectors: in 2024 => 13 months + 3 BC




# Pixel and HCAL phase 1 upgrades

Upgrade




New 4-layer pixel detector Pixel



## • Upgraded HCAL


- New photodetectors
- New electronics (frontend, backend)
- Improved longitudinal segmentation
- Improved background rejection, Missing  $E_T$  resolution and Particle Flow reconstruction



# Signal strength @3000 fb<sup>-1</sup>

**ATLAS** Simulation Preliminary


$\sqrt{s} = 14$  TeV:  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$




$$\mu = \sigma/\sigma_{\text{SM}}$$

**ATLAS** Simulation Preliminary




$\sqrt{s} = 14$  TeV:  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$



- With 3000 fb<sup>-1</sup> the couplings can be determined with high precision (a few %)



## VBF signal (2 jets)





# VV scattering: fully leptonic

Only background VV+jets, very low xsec

Number of events for  $20 \text{ fb}^{-1}$  (fully MC based, no systematics, 14 TeV)

CMS ZZ→4e, 4μ	N signal	N back.
500 GeV	2.2	1.9
>1 TeV	0.1	0.2



CMS ZW→μμμν	N signal	N back.
>1 TeV	0.9	0.8

ATLAS ZZ→2l2ν	N signal	N back.
500 GeV	6.4	3.0

ATLAS ZW→lllν	N signal	N back.
500 GeV	8	5
1.1 TeV	1.4	0.4

Example: ggF Higgs 300 GeV



Latest results:

B~ 6

S~ 10



• reso  $m_{4l}$  as expected

• improved reco-id efficiencies

(eg ele ID: TDR time 85-90% → today 95%)



## $pp \rightarrow ZZ + 2j \rightarrow 4\ell + 2j$ channel




$$\mathcal{L}_{\phi W} = \frac{c_{\phi W}}{\Lambda^2} \text{Tr}(W^{\mu\nu} W_{\mu\nu}) \phi^\dagger \phi$$

	300 $\text{fb}^{-1}$	3000 $\text{fb}^{-1}$
$c_{\phi W}/\Lambda^2$	34 $\text{TeV}^{-2}$	16 $\text{TeV}^{-2}$


Sensitivity to anomalous ZZ resonances in Vector boson scattering

## $pp \rightarrow WZ + 2j \rightarrow \ell' s + \nu + 2j$ channel

CMS Projection:  $\sqrt{s} = 14$  TeV,  $L = 300 \text{ fb}^{-1}$





CMS Projection:  $\sqrt{s} = 14$  TeV,  $L = 3000 \text{ fb}^{-1}$



Significance	$3\sigma$	$5\sigma$
SM EWK Scattering Discovery	$75 \text{ fb}^{-1}$	$185 \text{ fb}^{-1}$
$f_{T1}/\Lambda^4$ at $300 \text{ fb}^{-1}$	$0.8 \text{ TeV}^{-4}$	$1.0 \text{ TeV}^{-4}$
$f_{T1}/\Lambda^4$ at $3000 \text{ fb}^{-1}$	$0.45 \text{ TeV}^{-4}$	$0.55 \text{ TeV}^{-4}$

Sensitivity to anomalous WZ resonances in Vector boson scattering

## $pp \rightarrow ZZ + 2j \rightarrow 4\ell + 2j$ channel





model	$300 \text{ fb}^{-1}$	$3000 \text{ fb}^{-1}$
$m_{\text{resonance}} = 500 \text{ GeV}, g = 1.0$	$2.4\sigma$	$7.5\sigma$
$m_{\text{resonance}} = 1 \text{ TeV}, g = 1.75$	$1.7\sigma$	$5.5\sigma$
$m_{\text{resonance}} = 1 \text{ TeV}, g = 2.5$	$3.0\sigma$	$9.4\sigma$

Sensitivity to anomalous ZZ resonances in Vector boson scattering

# VV scattering: unitarity violation


**VV → VV**



**S channel**

**T channel**

**QGC**



Without the SM boson,  $W_L^+ W_L^- \rightarrow W_L^+ W_L^-$  violates unitarity at  $\sqrt{s} \geq 1.2$  TeV

W, Z masses ( $\rightarrow$  longitudinal degrees of freedom) arise from the BEH mechanism:

$$A(W_L^+ W_L^- \rightarrow W_L^+ W_L^-) \approx \frac{1}{v^2} \left( \boxed{-s - t} + \frac{s^2}{s - m_H^2} + \frac{t^2}{t - m_H^2} \right)$$

**VV scattering is the smoking gun for EWSB!**


Taken from “Prospects for VV scattering: latest news” by S. Bolognesi (JHU)  
talk at Implications of LHC results for TeV-Scale physics (March 2012)



# VV scattering as a probe for EWSB

## VV Scattering spectrum, $\sigma(VV \rightarrow VV)$ vs $M(VV)$

is the fundamental probe to test the nature of the BEH boson or to find an alternative EWSB mechanism




Search for possible resonances in VV scattering (VBS) spectrum




# VBS experimental signature

## Longitudinal plane




## Transverse plane



Signature: **forward-backward “spectator” jets with very high energy**

- ▶ *tagging jets (1): large  $p_T$ , large  $\Delta\eta$*
- ▶ *few jets between tagging jets*
- ▶ *final state  $\ell\nu\ell\nu$ :*
  - ▶ *leptons (2) between tagging jets*
  - ▶ *missing  $E_T(3)$*



**Six fermion  
final state**

# VBS final states

- According to the vector bosons' decays we have a multitude of possible final states. We can group them in:

- **Fully leptonic**

- $pp \rightarrow qq \ell\ell\ell\ell$  ( $\ell = \mu, e$ )
- $pp \rightarrow qq \ell\ell\ell\nu$
- $pp \rightarrow qq \ell\ell\nu\nu$

Clean

Can reconstruct  $m_{VV}$  (not with  $2\nu$ )

Very low yields...

- **Semi-leptonic**

- $pp \rightarrow qq \text{ jetjet } \ell\ell$
- $pp \rightarrow qq \text{ jetjet } \ell\nu$


Better yields...

Large backgrounds

## Detector needs

Excellent lepton ID, energy resolution, hermeticity, jet tagging at high  $\eta$


# VBS $2e2\mu$ candidate event






# WZ resonance

$pp \rightarrow WZ + 2j \rightarrow \ell + \nu + 2\ell + 2j$  channel



$$\mathcal{L}_{T,1} = \frac{f_{T1}}{\Lambda^4} \text{Tr}[\hat{W}_{\alpha\nu} \hat{W}^{\mu\beta}] \times \text{Tr}[\hat{W}_{\mu\beta} \hat{W}^{\alpha\nu}]$$




	300 $\text{fb}^{-1}$	3000 $\text{fb}^{-1}$
$f_{T1}/\Lambda^4$	1.3 $\text{TeV}^{-4}$	0.6 $\text{TeV}^{-4}$

Sensitivity to anomalous WZ resonances in Vector boson scattering




# WW resonance

$pp \rightarrow WW+2j \rightarrow 2\ell+2\nu+2j$  channel



$$\mathcal{L}_{S,0} = \frac{f_{S0}}{\Lambda^4} [(D_\mu \phi)^\dagger D_\nu \phi] \times [(D^\mu \phi)^\dagger D^\nu \phi]$$



model	300 $\text{fb}^{-1}$	3 $\text{ab}^{-1}$
$f_{S0}/\Lambda^4$	10 $\text{TeV}^{-4}$	4.5 $\text{TeV}^{-4}$


Sensitivity to anomalous WW resonances in Vector boson scattering

# Trigger challenge in 2012

Maintaining high trigger efficiency while keeping the trigger rate within budget was one of the biggest challenges of the CMS experiment in 2012


The experience obtained in 2012 with peak pileup of  $\sim 35$  events gives us confidence for high-luminosity running post Long Shutdown 1

## Trigger Cross-sections:




## HLT CPU time:

- linear with PU, no signs of runaway



# Tracking and b-tagging performance


Improvement of b-tagging efficiency  
with new pixel detector



b-tagging efficiency  $\sim 1.3x$  better  
2 b-jets  $\rightarrow (1.3)^2 \sim 1.69$

Primary vertex resolution improved by factor  $\sim 1.5 - 2$


Improvement in tracking efficiency w/  
new pixel detector, in ttbar events, as  
a function of pileup








# CMS Upgrade program

## LS1 and Phase 1





- Upgraded HCAL
  - New photodetectors
  - New electronics (frontend, backend)
  - Improved longitudinal segmentation
  - Improved background rejection, Missing  $E_T$  resolution and Particle Flow reconstruction
- Hadronic showers spread out with increasing depth



Reconstruction of hard collisions in high pileup environment requires detectors with very high granularity:



- efficient association of charged tracks to collision vertices
- reconstruction of charged and neutral particles in jets
- pileup neutrals corrected w/global energy density ( $\rho$ )

Physics with high pileup requires full particle flow reconstruction assuring:

- precise jet energy correction
- robust missing energy measurement
- efficient lepton isolation


Very efficient reconstruction code is needed to stay within computing budget

## Muon isolation



# Vector Boson Fusion (VBF)

Generic diagram for vector boson fusion (VBF) process



Signature: forward-backward  
“spectator” jets with very high energy

- Once the vector bosons decay, we have a **six-fermion** final state
  - The full set of  $qq \rightarrow 6$  fermions diagrams has to be considered
  - In order to investigate EWSB, one has to isolate VV processes from all other six-fermion final states
- ➡ Apply tight kinematic cuts

Typical kin. cuts

$$p_{T,j} > 20 \text{ GeV} \quad |\eta_j| < 5 \quad p_T^{\text{tag}} > 30 \text{ GeV} \quad |\eta_{j1} - \eta_{j2}| > 4.0$$

$$\eta_{j1} \cdot \eta_{j2} < 0 \quad m_{jj} > 600 \text{ GeV}$$

**Semileptonic is most promising: reasonable signal yield**


**Number of events for 20 fb<sup>-1</sup>** (fully MC based, no systematics, 14 TeV)

VV → Injj	ATLAS	N sign.	N back.	CMS	N sign.	N back.	ZV → lljj	CMS	N sign.	N back.
	500 GeV	6.2	16	500 GeV	337	20759		500 GeV	62	3415
	800 GeV	13	17							
	1.1 TeV	4.8	9.2	>1 TeV	45	3281		>1 TeV	5	348

For recent inclusive Higgs search:

- more sophisticated analysis developed (btag categories, angular analyses,  $m_{jj} = m_Z$  kinematic fit)
- data driven background

**Improved JES:**  $m_{jj}$  reso from 20-25% to 10-15%






# Ratios of partial widths @3000 fb<sup>-1</sup>

**ATLAS Preliminary (Simulation)**

$\sqrt{s} = 14$  TeV:  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$


$\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  extrapolated from 7+8 TeV



**ATLAS Preliminary (Simulation)**

$\sqrt{s} = 14$  TeV:  $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  ;  $\int \mathcal{L} dt = 3000 \text{ fb}^{-1}$

$\int \mathcal{L} dt = 300 \text{ fb}^{-1}$  extrapolated from 7+8 TeV



- With 3000 fb<sup>-1</sup> the couplings can be determined with high precision (a few %)

# Ratios of partial widths

## Scenario 1

CMS

partialWidths	300/fb (% err.)	3000/fb (% err)
r_bZ	24 / -18	12 / -9
r_gZ	16 / -13	8
r_tZ	18 / -15	9 / -7
r_WZ	15 / -12	7 / -6
r_topglu	32 / -24	17 / -13
r_Zglu	17 / -16	10 / -9
c_gluZ	12 / -11	8

Scenario 1: systematics as in 2012  
 Scenario 2: theory syst. scaled by a factor  $\frac{1}{2}$ , other systematics scaled by  $1/\sqrt{L}$

## Scenario 2

partialWidths	300/fb (% err.)	3000/fb (% err)
r_bZ	17 / -14	4.5
r_gZ	9	4.5
r_tZ	11	3.5
r_WZ	10 / -7	2.5
r_topglu	28 / -22	11
r_Zglu	11 / -10	5
c_gluZ	7.5 / -5.5	4

I would like to dedicate this talk to my father,  
**Prof. Giorgio Giacomelli**  
a worldwide known physicist,  
who passed away on January 30th 2014



**Giorgio Maria Giacomelli**  
**30/05/1931 - 30/01/2014**